

Analysis of $K\pi$ -scattering phase shift and existence of $\kappa(900)$ -particle

Shin ISHIDA, Muneyuki ISHIDA*, Taku ISHIDA**
Kunio TAKAMATSU***, and Tsuneaki TSURU**

*Atomic Energy Research Institute, College of Science and Technology,
Nihon University, Tokyo 101*

**Department of Physics, University of Tokyo, Tokyo 113*

***National Laboratory for High Energy Physics (KEK), Tsukuba 305*

****Department of Engineering, Miyazaki University, Miyazaki 988-12*

(Received February 1, 1997)

Recently we have shown an evidence for existence of σ -particle in the previous works; where the $\pi\pi$ S-wave phase shift is reanalyzed, by introducing a repulsive background suggested by the chiral symmetry, and by applying a new method of Interfering Breit-Wigner Amplitudes. In this work we also show, reanalyzing the $K\pi$ S-wave phase shift from a similar standpoint, an evidence for existence of $\kappa(900)$, possibly to be a member of σ -nonet.

§1. Introduction

In our previous works^{1), 2)} we have analyzed the iso-singlet S-wave $\pi\pi$ scattering phase shift, and have shown the existence of a resonance with mass of 535-650 MeV and width of about 350 MeV. These values are consistent with those of σ particle, the long-sought chiral partner of Nambu-Goldstone π meson, predicted by the linear σ model.³⁾ Independent analyses of the phase shift by several authors^{4), 5), 6)} have also suggested its existence. On the other hand anticipation for σ existence has been given recently with new interests both^{7) - 17)} theoretically and phenomenologically. As a matter of fact a low-mass isoscalar resonance, $f_0(400\sim 1200)$ or σ , has revived in the latest issue of Particle Data Group¹⁸⁾ after its missing over twenty years.

In the phase shift analysis^{1), 2)} on one hand, we have developed a new method of S-matrix parametrization in conformity with unitarity, Interfering Breit-Wigner Amplitude method, in which we use only a few parameters with direct physical meaning (*i.e.* masses and widths of resonances), in contrast with the conventional $\mathcal{K}$ -matrix method. On the other hand we have introduced an negative background phase δ^{BG} of hard core type phenomenologically. This type of background phase shifts¹⁹⁾ was observed historically in the α - α scattering, and also in the nucleon-nucleon scattering. In the relevant case of $\pi\pi$ system its origin seems to have some correspondence³⁾ to the “compensating” contact $\lambda\phi^4$ term required by the chiral symmetry in the linear σ model.

The σ -particle is a chiral partner of π -meson in the linear representation of chiral $SU(2)_L \times SU(2)_R$ group. Taking $SU(3)$ flavor symmetry into account, it is natural to expect existence of scalar σ -meson nonet as a chiral partner of pseudoscalar π -meson nonet. In the following we analyze the $I=1/2$ $K\pi$ scattering phase shift from

a similar standpoint to the $\pi\pi$ system, and actually show an evidence for existence of I=1/2 member of the σ -nonet, κ -meson.

§2. Applied formulas

We analyze the I=1/2 S-wave phase shift of $K\pi$ -scattering by Interfering Amplitude method in the case of three-channels ($K\pi$, $K\eta$ and $K\eta'$, denoted to 1, 2 and 3, respectively) with two-resonances (κ and $K_0^*(1430)$) from the $K\pi$ -threshold to $\sqrt{s} \sim 1.6$ GeV. ^{*)}

The relevant S-matrix element^{**) of $K\pi$ -scattering, S_{11} , is related to its phase shift (amplitude), $\delta_0^{\frac{1}{2}}(a_0^{\frac{1}{2}})$, by}

$$S_{11} = \eta_{11} e^{2i\delta_0^{\frac{1}{2}}} = 1 + 2ia_0^{\frac{1}{2}}, \quad (1)$$

where η_{11} is the elasticity. S_{11} is given by product of “individual” resonance-S-matrices $S_{11}^{(R)} (R = \kappa, K_0^*)$

$$S_{11} = e^{2i\delta^{BG}} \prod_{R=\kappa, K_0^*} S_{11}^{(R)}; \quad (2)$$

The unitarity of “total” S-matrix (2) is now easily seen to be satisfied by the unitarity of individual S matrices. In Eq.(2) we have also introduced a negative background phase δ^{BG} , taken as a hard core type phenomenologically,

$$\delta^{BG} = -|\mathbf{p}_1| r_{c0}^{\frac{1}{2}}; \quad |\mathbf{p}_1| = \frac{\sqrt{(s - m_\pi^2 - m_K^2)^2 - 4m_\pi^2 m_K^2}}{2\sqrt{s}}, \quad (3)$$

$|\mathbf{p}_1|$ being the CM momentum of the πK system.

Each of $S_{11}^{(R)}$ is given by a corresponding amplitude $a_{11}^{(R)}$ taken as a simple relativistic Breit-Wigner form

$$S_{11}^{(R)} = 1 + 2ia_{11}^{(R)}, \quad a_{11}^{(R)} = \frac{-\sqrt{s}\Gamma_R^1(s)}{s - M_R^2 + i\sqrt{s}\Gamma_R^{\text{tot}}(s)}, \quad (4)$$

where $\Gamma_R^{\text{tot}}(s)(\Gamma_R^1(s))$ is a total width (partial width of channel 1) of the resonance R , given by

$$\Gamma_R^{\text{tot}}(s) = \sum_{i=1}^3 \Gamma_R^i(s); \quad \Gamma_R^i(s) = \frac{\rho_i}{\sqrt{s}} g_{Ri}^2 = g_{Ri}^2 |\mathbf{p}_i| / 8\pi s \quad (i = 1, 2, 3). \quad (5)$$

Here g_{Ri} ’s are coupling constants to the channel i of resonance R , and the CM momentum $|\mathbf{p}_i|$ for $i = 2, 3$ are defined in a similar way as in Eq.(3). Thus parameters to be used for the fit are totally nine, *i.e.* resonance masses M_R ’s ($R=\kappa, K_0^*$), their coupling constants g_{Ri} ’s ($i=1,2,3$), and repulsive core radius $r_{c0}^{\frac{1}{2}}$.

^{*)} Contributions from the other channels such as $K\pi\sigma$ and $K\pi\pi\pi$ are expected to be suppressed by a phase space factor.

^{**) The other elements of $S_{ij} (i \neq 1 \text{ and/or } j \neq 1)$ are now irrelevant since of present experimental situations in the corresponding processes.}

§3. Mass and width of κ and core radius

A high statistics data of the reaction $K^-p \rightarrow K^-\pi^+n$ was obtained with 11 GeV/c beam using LASS spectrometer at SLAC.²⁰⁾ Spherical harmonic moments were used to perform an energy independent Partial Wave Analysis of the $K^-\pi^+$ system from threshold to 2.6 GeV, with t -dependent parametrization of the production amplitudes. The obtained $K^-\pi^+$ -scattering amplitudes are the sum of $I=1/2$ and $3/2$ components. The $I=1/2$ S-wave amplitude $a_0^{\frac{1}{2}}$ was determined by subtracting the $I=3/2$ ($K^+\pi^+/K^-\pi^-$) component, obtained independently by another experiment at SLAC.²¹⁾ Here, the “overall” phase was fixed by imposing elasticity constraint to the amplitude in the $m_{K\pi}$ region below 1.29 GeV. We use this amplitude between $K\pi$ threshold and 1.6 GeV for the analysis.

Figure 1(a) and (b) show the result of the best fit to $\delta_0^{\frac{1}{2}}$ and $|a_0^{\frac{1}{2}}|$, respectively, by solid line. The obtained parameters are collected in Table I. The most remarkable feature is that we identify a low-mass resonance κ with mass of about 900 MeV in the slowly-increasing phases between the threshold and 1300 MeV. This is due to the role of “compensating” repulsive background δ_{BG} , whose existence is necessarily required from Chiral Symmetry (see ii) and iii) of the following supplementary discussions). As a matter of fact, the original LASS analysis of the data²⁰⁾, where a positive δ_{BG} with an effective range formula was introduced, led to existence of only one state $K_0^*(1430)$ with high mass. Since of the compensation between contributions due to κ and the repulsive core, the mass value of κ (M_κ) and its coupling to the $K\pi$ channel ($g_{\kappa 1}$) are correlated to the core radius r_c . To clarify this situation, various fits are performed with a series of fixed r_c values between 0 to 5.5 GeV⁻¹. Fig. 1(c) shows the values of χ^2 , M_κ , $g_{\kappa 1}$, $M_{K_0^*}$, and $g_{K_0^* 1}$ as functions of r_c . M_κ and $g_{\kappa 1}$ decrease as r_c becomes larger, while $M_{K_0^*}$ and $g_{K_0^* 1}$ do not show such correlations because these values are constrained mainly by the steep phase increase around 1.4 GeV. In the range of 2~5 GeV⁻¹, the χ^2 value shows a parabolic shape, and makes its minimum at $r_c=3.57$ GeV⁻¹ where we get the best fit given in Fig. 1(a) and (b) ($\chi^2 = 57.0$ for 42 degrees of freedom; 51 data points with 9 parameters). When r_c becomes smaller than 2 GeV⁻¹, the values of M_κ and $g_{\kappa 1}$ increase steeply, and the contribution of “ κ -meson resonance” has no more meaning than the positive background. A fit with r_c setting to zero gives large M_κ and $g_{\kappa 1}$ values (6.4 GeV and 39 GeV, respectively), and becomes essentially similar to the LASS analysis. This fit has the χ^2 value of 96, which is larger by 40 than of our best fit. *)

From the χ^2 behavior in Fig. 1(c), we can obtain the upper and lower bounds of error of M_κ , $g_{\kappa 1}$ and r_c which are given in Table I as five standard deviations from the best fit (+25 χ -squares). Corresponding curves with upper and lower values of relevant parameters are also shown in Fig. 1(a) and (b), respectively, by dotted ($r_c=3.1$ GeV⁻¹) and dashed ($r_c=3.975$ GeV⁻¹) lines.

*) In Ref. 20) almost all data points are given with no accurate errors. We will regard the original errors of $K^-\pi^+$ amplitude data²²⁾ equivalent to our relevant errors of $I=1/2$ scattering amplitude, which might be smaller than those of 20). This may be a reason why we get a χ -square value larger than that in the LASS analysis.




Fig. 1. Fits to $I=1/2$ $K\pi$ S-wave scattering amplitude; (a) phase shift $\delta_0^{1/2}$, and (b) magnitude of amplitude $|a_0^{1/2}|$. The solid lines are the best fit with $r_c=3.57\text{GeV}^{-1}$, while the dotted and dashed lines are fits with $r_c=3.1$ and 3.975GeV^{-1} , respectively. (c) χ^2 , M_κ , g_κ , $M_{K_0^*}$, and $g_{K_0^*}$ behaviors as functions of core radius r_c . Vertical lines represent $r_c=3.57$, 3.1, and 3.975 GeV^{-1} , corresponding to the best fit and the fit with ± 5 sigma deviations.

Table I. Resonance parameters of $\kappa(900)$, $K_0^*(1430)$ and core radius. The errors correspond to five standard deviations from the best fit. Two kinds of width, $\Gamma^{(p)}$ and $\Gamma^{(d)}$ defined as $\Gamma^{(p)} = \Gamma_R^i(s = M^2)$ (Eq.(5)), $\Gamma^{(d)} = N^{-1} \int ds \Gamma(s) / [(s - M^2)^2 + s\Gamma(s)^2]$; $N = ds \int 1 / [(s - M^2)^2 + s\Gamma(s)^2]$, considering broadness of relevant widths.

	M_κ	$g_{K\pi}$	$\Gamma_{K\pi}^{(p)}$	$\Gamma_{K\pi}^{(d)}$
$\kappa(900)$	$905_{-30}^{+65} \text{ MeV}$	$6150_{-650}^{+1200} \text{ MeV}$	$545_{-110}^{+235} \text{ MeV}$	$470_{-90}^{+185} \text{ MeV}$
$K_0^*(1430)$	$1410_{-15}^{+10} \text{ MeV}$	$4250_{-70}^{+380} \text{ MeV}$	$220_{-5}^{+40} \text{ MeV}$	$220_{-5}^{+40} \text{ MeV}$

$r_{c0}^{1/2}$
$3.57_{-0.45}^{+0.40} \text{ GeV}^{-1} \text{ (} 0.70_{+0.08} \text{ fm)}$




Fig. 2. $I=3/2$ $K\pi$ scattering phase shift. Fitting by hard core formula is also shown.

Table II. Phenomenological core radii r_{c0}^I in $\pi\pi$ and $K\pi$ systems

	$(\pi\pi)_{I=0}$	$(K\pi)_{I=1/2}$	$(\pi\pi)_{I=2}$	$(K\pi)_{I=3/2}$
r_c	$0.60 \pm 0.07 \text{ fm}$	$0.70 \pm 0.09 \text{ fm}$	0.17 fm	0.16 fm

The respective coupling constants to the 2nd channel $K\eta$ are obtained to be much smaller than those to the $K\pi$ channel, *i.e.*, $g_{\kappa 2} \lesssim 1.0 \text{ GeV}$ and $g_{K_0^* 2} \lesssim 0.9 \text{ GeV}$, which are consistent to the elasticity constraint mentioned above. The $g_{\kappa 3}$ and $g_{K_0^* 3}$, couplings to $K\eta'$ channel, are obtained with much larger uncertainties, and their values are omitted here.

§4. Repulsive background in $K\pi$ - and $\pi\pi$ -systems

In the present analysis, leading to the existence of κ -meson, introduction of a negative background phase δ^{BG} of hard core type plays an essential role. This is a similar situation as for σ -existence in $\pi\pi$ -scattering. In the $I=2$ $\pi\pi$ system, there are no known and/or expected resonances, and this repulsive type phase shift itself is, if it exists, expected to be observed directly. Actually a good fit to the experimental data was obtained²⁾ by a similar formula of hard core type as Eq.(3) with the core radius $r_{c0}^2 = 0.17 \text{ fm}$. The same situation is expected in the $I=3/2$ $K\pi$ scattering, and we have made an similar analysis on the relevant $K\pi$ -scattering phase shift.²¹⁾ The result is given in Fig. 2. The best fit is obtained with the core radius $r_{c0}^{3/2} = 0.16 \text{ fm}$, although the fit is somewhat worse than in the case of $I=2$ $\pi\pi$ system.

The values of phenomenological core radii in the $\pi\pi$ and $K\pi$ systems are collected in Table II. It is quite interesting that they are almost same within the non-exotic channels (that is, $r^0=r^{1/2}$) and within the exotic channels ($r^2=r^{3/2}$), respectively. It seems to be reasonable from the viewpoint of $SU(3)$ flavor symmetry.

§5. Supplementary discussions

Here we give some additional comments on the results of our analysis:

i) It may be interesting and important to compare the properties of κ -meson obtained above with predictions of the various theoretical models. The SU(3) Linear σ Model with the $U_A(1)$ breaking term(L σ M1)^{23), 24), 25)}, the SU(3) Linear σ Model without it(L σ M2)²⁶⁾, and the extended Nambu Jona-Lasinio type model (ENJLM)^{27), 28)} (including also the $U_A(1)$ breaking term) give the κ -masses, respectively, as $m_\kappa = 1.2$ ²⁴⁾, 0.8 ²⁶⁾ and $0.4 \sim 0.9$ ^{27) *} in GeV, which have a large uncertainty. The values of κ decay width are given to be $1.2(0.1)$ GeV in L σ M1 (L σ M2). The properties of κ meson in Table I seem not inconsistent with those predicted by the SU(3)-theoretical models with the $U_A(1)$ breaking term(L σ M1 and ENJLM). Accordingly our κ meson can be regarded as the member of σ -nonet, although further investigations are necessary.

ii) Concerning a possible origin of the repulsive background phase δ_{BG} , introduced phenomenologically in this analysis, we should like to note its similarity³⁾ to the $\lambda\phi^4$ term in L σ M. The $\lambda\phi^4$ term represents a strong repulsive and contact (zero-range) interaction between pions and seems to have a plausible property as an origin of δ_{BG} , at least, in the low energy region, where the structures of composite pions may be neglected.

In $K\pi(\pi\pi)$ -scattering in L σ M a contribution due to intermediate $\kappa(\sigma)$ -production in all s, t, u -channels almost cancels^{3), 5)} in the low energy region with a repulsive force from the $\lambda\phi^4$ interaction. This leads effectively to the derivative (thus small) coupling⁶⁾ of Nambu-Goldstone boson. This cancellation mechanism is guaranteed by chiral symmetry and PCAC. It is notable that in the usual Breit-Wigner formula of S-wave resonance, a non-derivative coupling of κ and σ resonance is supposed, without taking the “compensating” repulsive interaction into account. This seems to be a reason why σ and κ resonances have been overlooked in the many phase shift analyses thus far made.

The result given in Table II, that the repulsive core radii in non-exotic channels are much larger than those in exotic channels, may be given²⁹⁾ some reason in L σ M as follows: In exotic channels a large amount of strong repulsive force due to $\lambda\phi^4$ interaction is canceled by the attractive force due to crossed-channel exchange of relevant scalar-mesons, while in non-exotic channels there remains some amount of the repulsive force (going to compensate the attractive force due to s -channel intermediate production of the scalar mesons in the threshold).

iii) In analysis of the $\pi\pi(K\pi)$ -scattering, the importance of ρ (K^*) meson effects is often pointed out.³⁰⁾ In the S-wave scattering these vector mesons contribute only through the crossed channel exchange diagrams, which are necessarily accompanied by the “compensating” derivative ϕ^4 interaction,³¹⁾ similarly in the case of the $\lambda\phi^4$ interaction to the κ (or σ) exchange. They exactly cancel with each other at the

*) This value is quoted in their analysis of case A, where $m_\sigma = 0.604\text{GeV}$, close to our value^{1), 2)}.

Table III. Candidate for Chiralons; members of chiral scalar-meson nonet.

		$M(\text{MeV})$	$g(\text{MeV})$	$\Gamma^{(p)}(\text{MeV})$	
I=0($n\bar{n}$)	$\sigma(535\text{-}650)$	585 ± 20	3600 ± 350	385 ± 70	f_{1c}
I=0($s\bar{s}$)	σ'	?	?	?	f'_{1c}
I=1/2	$\kappa(900)$	905 ± 50	6150 ± 900	545 ± 170	K_{1c}
I=1	δ	?	?	?	a_{1c}

threshold and give only small effects in the low energy region, which may be regarded as included in the background phase.

iv) Finally we give a comment on the behavior of the background phase. In I=2 (I=3/2) channel of S-wave $\pi\pi(K\pi)$ -scattering the fit of phase shifts by hard core formula is satisfactory below $\sqrt{s} \sim 1.4$ GeV, as was shown in Fig. 1 in Ref. 2) (Fig. 2). However, the experimental phase shifts in the higher energy region seem to be decreasing³²⁾. This is a very interesting phenomenon, which reminds us of the soft core in nucleon-nucleon scattering. In this work we applied the hard core type background, implicitly supposing “local” π and K mesons. The above mentioned “soft core” type behavior of phase shifts in the comparatively high energy region seems to suggest the composite structure of π and K as $q\bar{q}$ -bound states.

§6. Concluding remarks

We have shown a strong evidence for existence of σ -meson in the previous work^{1),2)} and of κ -meson in the present work. The existence of these particles has a significant importance in hadron spectroscopy. Since of their light masses (and, for σ , of its vacuum quantum number), they will appear in various processes such as $K \rightarrow 2\pi$ decay¹⁰⁾, K_{l4} decay¹³⁾, etc..

We have argued that κ meson observed in the present analysis is a member of σ -meson nonet, chiral partner of π -meson nonet. As was discussed in Ref.1), these particles (we call them “Chiralons”) should be regarded as being different from the ordinary P-wave excited states of $q\bar{q}$ system since of their light masses. This discrimination may have also some theoretical reasons: In the extended Nambu Jona-Lasinio model as a low energy effective theory of QCD (, in which the existence of σ -meson is predicted), only local composite quark and anti-quark operators are treated, thus missing L -excited states in principle. The present status of “Chiralons,”¹⁾ is summarized in Table III.

Classification of low mass scalar mesons is still in confusion. One of its main reasons seems to come from mis-identification of the chiral σ -nonet with the $q\bar{q}$ 3P_0 nonet. The properties of this extra-nonnet should be further investigated also through the many other production processes, such as pp -central collision¹⁷⁾, Υ and Ψ decays. In this connection especially the properties of observed resonances, $f_0(980)$ and $a_0(980)$, are to be clarified in relation^{24),26)} to the other members of chiralons, σ' with I=0 and δ with I=1.

Acknowledgements

The authors should like to express their deep gratitude to S. Suzuki (Nagoya university) who has kindly informed us of Ref. 22).

References

- [1] S. Ishida, M.Y. Ishida, H. Takahashi, T. Ishida, K. Takamatsu and T. Tsuru, Prog. Theor. Phys. **95** (1996), 745.
- [2] S. Ishida, T. Ishida, M.Y. Ishida, K. Takamatsu, T. Tsuru and H. Takahashi, Nihon Univ. preprint NUP-A-96-11, KEK preprint 96-131 (hep-ph/9610359).
- [3] M.Y. Ishida, Prog. Theor. Phys. **96** (1996), 853.
- [4] R. Kamiński, L. Leśniak and J.-P. Maillet, Phys. Rev. **D50** (1994), 3145, private communication.
- [5] N.A. Törnqvist and M. Roos, Phys. Rev. Lett. **76** (1996), 1575; N.A. Törnqvist, Z. Phys. **C68** (1995), 647.
- [6] M. Harada, F. Sannino and J. Schechter, Phys. Rev. **D54** (1996), 1991.
- [7] R. Delbourgo and M.D. Scadron, Phys. Rev. Lett. **48** (1982), 379.
- [8] T. Hatsuda and T. Kunihiro, Prog. Theor. Phys. **74** (1985), 765; *Phys. Rep.* **247** (1994), 221.
- [9] T. Kunihiro, Prog. Theor. Phys. Suppl. **120** (1995), 75.
- [10] E.P. Shabalin, Yad.Fiz. **48** (1988), 272 (Sov.J.Nucl.Phys. **48** (1988), 172). T. Morozumi, C.S. Lim and A.I. Sanda, Phys. Rev. Lett. **65** (1990), 404. M. Takizawa, T. Inoue and M. Oka, Prog.Theor.Phys.Suppl. **120** (1995), 335.
- [11] G. Mennessier, Z. Phys. **C16** (1983), 241; S. Minami, Prog. Theor. Phys. **81** (1989), 1064.
- [12] E. Beveren, T.A. Rijken, K. Metzger, C. Dullemond, G. Rupp and J.E. Ribeiro, Z. Phys. **C30** (1986), 615.
- [13] E.P. Shabalin, Yad.Fiz. **49** (1989), 588 (Sov.J.Nucl.Phys. **49** (1989), 365).
- [14] H. Yoshino *et al.*, Prog. Theor. Phys. **95** (1996), 353.
- [15] M. Svec, Phys. Rev. **D53** (1996), 2343.
- [16] H. Shimizu, contributed paper on Particle and Nuclei XIII International Conference - PANIC XIII, Perugia, Italy, June 1993.
- [17] T. Ishida *et al.*, in: Proc. sixth International Conference on HADRON Spectroscopy (Manchester, July'95), ed. M.C.Birse, G.D.Lafferty and J.A.McGovern (World Scientific) p.451 (KEK-Preprint 95-159 (1995)). T. Ishida, Doctor Thesis of Tokyo University(1996). D. Alde *et al.*, to be published in Physics Letters B (KEK Preprint 96-174).
- [18] Review of Particle Properties, Phys. Rev. **D54** Part I (1996) p.329 and also p.355.
- [19] M. Taketani *et al.*, Prog. Theor. Phys. Suppl. No.39(1967); No.42(1968). In particular see Chapter 7 (S. Otsuki, No.42, p.39) and also Chapter 6 (N. Hoshizaki, No.42, p.1).
- [20] D. Aston *et al.*, Nucl. Phys. **B296** (1988), 493.
- [21] P. Estabrooks *et al.*, Nucl. Phys. **B133** (1978), 490.
- [22] N. Awaji, Doctor thesis of Nagoya university (1986).
- [23] G. Gasiorowicz and D.A. Geffen, Rev.Mod.Phys. **3** (1969), 531.
- [24] E.P. Shabalin, Yad.Fiz. **41** (1985), 260 (Sov.J.Nucl.Phys. **42** (1985), 164).
- [25] T. Morozumi, A.I. Sanda and A. Soni, Phys. Rev. **D46** (1992), 2240.
- [26] M.D. Scadron, Phys. Rev. **D26** (1982), 239.
- [27] M. Takizawa, K. Tsushima, Y. Kohyama and K. Kubodera, Nucl. Phys. **A507** (1990), 611.
- [28] S. Klimt, M. Lutz, U. Vogl and W. Weise, Nucl. Phys. **A516** (1990), 429.
- [29] M.Y. Ishida, Proc. of YITP conference, 1996, kyoto, Yukawa Hall, to be published in Soryuusiron-kenkyuu; Proc. of the Workshop on Particle Physics at K arena with 50-GeV PS, KEK, Tsukuba, Japan Dec 19-20,1996, edited by N.Sasao, S.Sugimoto, Y.Kuno and T.Komatsubara, Inst. for Nuclear Study, Univ. of Tokyo.
- [30] B.S. Zou and D.V. Bugg, Phys. Rev. **D50** (1994), 591. N. Isgur and J. Speth, Phys. Rev. Lett. **77** (1996), 2332 and the references therein.
- [31] S. Weinberg, Phys.Rev. **166** (1968), 1568. T. Shiozaki, Prog. Theor. Phys. Suppl. Extra Number (1968) 44. M. Bando, T. Kugo and K. Yamawaki, *Phys.Rep.* **164** (1988), 217.

- [32] N.B. Durusoy *et al.*, Phys. Lett. **45B** (1973), 517.